

Protocol 700-M4

Nausea and Vomiting

Rev: 2/18

BLS Treatment

- ❖ Treat life threats. (See Procedure 701 *Life Threats*)
- ❖ Pay particular attention to maintaining a patent airway, and protecting the patient from aspiration.
- ❖ Consider underlying causes for nausea/vomiting, and treat as appropriate
- ❖ Attempt non-invasive methods of reducing nausea/vomiting, including reducing environmental stimulation, providing fresh air, applying oxygen, reducing unpleasant odors, and using distracting techniques.
- ❖ Prepare for transport/transfer of care.

ALS Treatment

- ❖ Treat life threats. (See Procedure 701 *Life Threats*)
- ❖ **Ondansetron**
 - 4 mg IV/IO/IM/ODT.
 - May repeat every 5-10 minutes as needed Max 16 mg.
- ❖ Transport/ Contact Base Station as needed.

Special Considerations

- ❖ **Ondansetron** is considered safe for pregnancy (Class B)
- ❖ **Ondansetron** rarely causes sedation, and is typically well tolerated by all ages of patients.
- ❖ Nausea/vomiting is a symptom. Be aware of underlying causes*
- ❖ **Ondansetron** is contraindicated in patients with diagnosed Long QT Syndrome, and for those who are currently taking **Amiodarone**, Haldol, Methadone, Procainamide, or Seroquel.

*Causes of Nausea	
Narcotics	Toxic Ingestion
Motion Sickness	Gastroenteritis
Head Injury	Acute MI
Abdominal Pain	Stroke
Pregnancy	



David Ghilarducci MD
EMS Medical Director