

BLS Treatment

- ❖ Ensure scene safety
- ❖ Remove the victim from the source of exposure
- ❖ Treat life threats. (See Procedure 701 *Life Threats*)
- ❖ Administer high flow oxygen via a NRB
- ❖ Evaluate the patient for facial burns, hoarseness, black sputum, and soot in the nose and/or mouth
- ❖ Completely remove the victim's clothing prior to transport.
- ❖ Perform spinal immobilization if c-spine precautions are indicated
- ❖ Assess and treat for thermal and/or traumatic injuries (See Policy *E4 Burns or Policy T1 Trauma*)
- ❖ Manage the patient's airway early. Use BVM with airway adjuncts as appropriate
- ❖ Treat bronchospasms and airway problems as necessary (See Policy *R1 Respiratory Distress*)
- ❖ Place patient in position of comfort.
- ❖ Observe for signs of severe respiratory distress (Table 1)
- ❖ Prepare for transport/transfer of care.

Table 1: Signs of Severe Respiratory Distress

• ALOC	• low SpO ₂ ,
• Sig. accessory muscle use	• poor skin signs
• fatigue	• Elevated EtCO ₂
	• inability to speak

ALS Treatment

- ❖ Manage the patient's airway early. Intubate the patient if necessary (See Procedure 704, *Advanced Airway Management*)
- ❖ Consider a **Normal Saline** bolus
- ❖ Transport/Contact Base Station.

Special Considerations

- ❖ **Warning:** Pulse oximetry values may be unreliable in smoke inhalation patients.
- ❖ Cyanide and/or the combination of cyanide and carbon monoxide may be responsible for the majority of smoke inhalation deaths

David Ghilarducci MD