

Protocol 700-N2-P

Seizure

Rev: 2/18

BLS Treatment

- ❖ Treat life threats. (See Procedure 701 *Life Threats*)
- ❖ Protect patient from injury. Spinal precautions as indicated
- ❖ Check for Hypoglycemia
 - Perform Blood Glucose check
 - if less than 60 mg/dl treat as needed.
 - If conscious and patient can swallow on command, administer glucose paste or let patient self-administer glucose product.
 - If unconscious, place a dime size amount of glucose paste under the tongue.
- ❖ Prepare for transport / transfer of care.

ALS Treatment

- ❖ Treat life threats. (See Procedure 701 *Life Threats*)
- ❖ Do BG Chem and if less than 60 dl/ml proceed with **Dextrose**.
- ❖ If persistent seizures, administer **Midazolam** 0.05 mg/kg IV/IO to a maximum of 2 mg total, or 0.1mg/kg IM to a maximum of 5 mg total.
 - After max dose, contact Base Station for additional doses. In higher doses **Midazolam** may cause respiratory depression.
- ❖ Transport
- ❖ Contact Base Station.

Special Considerations

- ❖ Status epilepticus is a true medical emergency defined as either continuous seizure lasting at least five minutes or two or more discrete seizures between which there is an incomplete recovery of consciousness.
- ❖ Continuous capnography, pulse oximetry, and blood pressure monitoring are mandatory during and after administration of **Midazolam**.



David Ghilarducci MD
EMS Medical Director